

IN THE CLAIMS

Please amend the claims as follows:

Claim 1 (Original): A process for preparing prepolymers containing isocyanate groups by reacting

- a) diisocyanates with
- b) compounds having at least two hydrogen atoms which are reactive toward isocyanate groups in the presence of
- c) catalysts, and subsequently separating off the excess monomeric diisocyanates,

wherein

the diisocyanates a) used are unsymmetrical diisocyanates and the catalysts c) used are organometallic catalysts and these organometallic catalysts are removed, blocked or deactivated before the monomeric diisocyanates are separated off.

Claim 2 (Original): A process according to claim 1, wherein unsymmetrical diisocyanates used are tolylene 2,4'-diisocyanate, diphenylmethane 2,4'-diisocyanate and/or isophorone diisocyanate.

Claim 3 (Original): A process according to claim 1, wherein the unsymmetrical diisocyanate used is diphenylmethane 2,4'-diisocyanate.

Claim 4 (Original): A process according to claim 1, wherein the metal catalysts are selected from the group consisting of organometallic compounds of the metals of groups IVA, VA, IVB, VB and VIIB.

Claim 5 (Original): A process according to claim 4, wherein the metal catalysts contain ligands.

Claim 6 (Original): A process according to claim 4, wherein the ligands used are carboxylate anions, alkoxides, enolates, thiolates, mercaptides and alkyl ligands and combinations thereof.

Claim 7 (Original): A process according to claim 4, wherein the ligands are used in the form of chelating systems.

Claim 8 (Original): A process according to claim 1, wherein the metal catalysts are selected from the group consisting of dimethyltin, dibutyltin and dioctyltin dilaurate, bis(dodecylmercaptide), bis(2-ethylhexylthioglycolate), diacetate, maleate, bithioglycerol; octyltin tris(2-ethylhexylthioglycolate), bis( $\beta$ -methoxycarbonyl-ethyl)tin dilaurate, tetraisopropyl titanate, tetra-tert-butyl orthotitanate, tetra(2-ethylhexyl)titanium and bis(ethylacetoacetato)titanium diisopropoxide, bismuth(III) tris(2-ethylhexanoate) and bismuth laurate.


Claim 9 (Original): A process according to claim 1, wherein the metal catalysts are homogeneous catalysts.

Claim 10 (Original): A process according to claim 1, wherein the metal catalysts are heterogeneous catalysts.

Claim 11 (Original): A process according to claim 1, wherein the metal catalysts have been applied to supports.

Claim 12 (Original): A process according to claim 1, wherein the organometallic catalysts are deactivated by means of Lewis-acid metal deactivators.

Claim 13 (Original): A process according to claim 1, wherein the organometallic catalysts are deactivated by means of compounds of the general formula (I)



where  $R_1$  and  $R_4$  are, independently of one another, any organic radicals such as a linear, branched or cyclic alkyl radical, a linear, branched or cyclic alkenyl radical, a linear, branched or cyclic hydroxy, halogen, amino or thioalkyl radical,  $R_2$  and  $R_3$  are each, independently of one another, either nothing or hydrogen,  $X_1$  and  $X_4$  are each, independently of one another, either nothing or oxygen,  $X_2$  and  $X_3$  are Lewis-acid substituents, for example a halogen, O, OH,  $NH_2$ ,  $NO_2$ , SH and  $n$  is an integer from 1 to 5.

Claim 14 (Original): A process according to claim 13, wherein the compounds of the general formula (I) are organic carboxylic acids which are functionalized on the  $\beta$ -carbon atom (C3) relative to the carbon atom (C1) of the acid group ( $-C(1)OOH$ ) from the group consisting of  $\beta$ -hydroxycarboxylic acids,  $\beta$ -aminocarboxylic acids,  $\beta$ -ketocarboxylic acids and 1,3-dicarboxylic acids and their esters.

Claim 15 (Original): A process according to claim 13, wherein the compounds of the general formula (I) are selected from the group consisting of citric acid, malic acid, tartaric acid, acetoacetic acid, 2-chloroacetoacetic acid, benzoylacetic acid, acetonedicarboxylic acid,

dehydroacetic acid, 3-oxovaleric acid and malonic acid and also the associated esters in each case.

Claim 16 (Currently Amended): A prepolymer which contains isocyanate groups and has a content of monomeric diisocyanates of from 0.01 to 0.5% by weight, based on the weight of the prepolymer, and a content of ABA structures of at least 80% and can be prepared according to ~~any of claims 1 to 13~~ claim 1.

Claim 17 (Currently Amended): The ~~use of~~ method of using compounds of the general formula (I) of claim 13 for the deactivation of organometallic catalysts in the preparation of prepolymers containing isocyanate groups.